


# *Determining lamprey species composition, larval distribution and adult abundance in the Deschutes River sub-basin*



*Jennifer Graham  
Confederated Tribes of Warm Springs Reservation, Oregon  
Department of Natural Resources*



*Project  
Area*





# *Objectives*

---

*Objective 1: Determine larval distribution and associated habitat*

*Objective 2: Estimate the number of lamprey out-migrants by developmental stage*

*Objective 3: Evaluate the feasibility of estimating the escapement of adult lamprey and determine harvest rates at Sherar's Falls.*





*Larval Distribution and Associated Habitat*

# *Methods*

---

*~ Randomly selected 1 sample site per 10 Rkm*



*~ May - September*



*~ Three-tiered sampling methodology*






# Methods – Habitat & Water Quality


## Tier 2 – Transect

Wetted Channel Width  
Bankfull Channel Width  
Canopy Density



## Tier 1 – Reach

Conductivity  
pH  
Water Temperature  
Flood Prone Width  
Channel Slope



## Tier 3 – Sub-sample

Mean Water Depth  
Water Velocity  
Substrate Composition  
Habitat Type



# *Methods – Lamprey sampling*

## *Tier 3 – Sub-sample*




- ~ Backpack electrofisher (AbP-2)
- ~ 2, 90-s samples
- ~ Lamprey
- ~ Anesthetized, identified, weighed and measured
- ~ Returned to collection area



# Results - Streams

~ 384 m<sup>2</sup> sampled

~ 4 of 13 streams  
sampled contained  
ammocoetes





# *Results – Lamprey*

*~ 131 ammocoetes collected*



*~ Total length range: 25 mm – 145 mm*

*~ Mean total Length: 77 mm*





*Regression*

---

*P-Value*

*Lamprey presence vs. Woody debris*

*0.000*

*Lamprey presence vs. Depositional Area*

*0.011*

*However, sample sizes were small!!*





# *Plans for 2004*

---

- ~ Larval distribution and habitat data collection*
  - ~ Remainder of perennial streams in the lower Deschutes*
  - ~ Mainstem Deschutes*





*Estimate lamprey out-migrants and  
out-migration timing*




# Methods

- ~ Fished 2 screw traps
- ~ 5 days/wk; 24 hrs/day




- ~ Collected lamprey
- ~ Anesthetized
- ~ Identified to species
- ~ Developmental stage
- ~ Length

# Results – Warm Springs Timing






# Results – Warm Springs Timing



# Results – Shitike Creek Timing



# Trap efficiencies

- ~ *Ammocoetes captured through electrofishing*
- ~ *Marked with elastomer*
- ~ *Placed in holding box*
  - ~ *Checked after 24 hours*



- ~ *Multiple trap holding efficiencies*
  - ~ *0% holding efficiency*
- ~ *No out-migrant estimates completed*





# *Plans for 2004*

*~ Continue collecting timing data*

*~ Modify existing traps*

*~ Conduct trap efficiencies*

*~ Mark-recapture to estimate out-migrant numbers*





# *Tribal Harvest and Escapement Estimates*




*Picture courtesy of Lyman Jim*



# *Sherar's Falls*

## DESCHUTES RIVER BASIN





# Methods – Mark-recapture

## “Lamprey pots” (USFWS)

- ~ Late-June – early-July
- ~ Multiple modifications
- ~ Fished various locations in fish ladder



## Long-handled dip net

- ~ Late-June – August
- ~ Fished each pool 1 time/hr
- ~ Same location each time



## *Results - Tagging*

*Fin Clip*

*Floy Tag*

*199 Lamprey Tagged*



# Results – Recapture

*35 Recaptures (17.6% recapture rate)*

*11 Netting; 22 Creel; 2 Tribal member returns*



*Primary Tag Retention 77%*





# Methods – Tribal Harvest


- ~ Access site creel survey
  - ~ Mid-June - August
  - ~ 4 weekdays; 1 weekend day
  - ~ 1 hr. after sunset – 3 am
- ~ Creel lamprey
  - ~ Examined for fin clip and floy tag
  - ~ Total length measured
- ~ Creel data expanded to estimate total tribal harvest



# *Results – Tribal Harvest*

- ~ 21 interviews conducted*
- ~ 614 lamprey collected*
- ~ 1,017 estimated harvest*
  - ~ 9.25 lamprey/hr*
  - ~ Mean length: 62 cm*
  - ~ Range: 50-74 cm*

*Roger Jim, Sr.  
Celilo Falls*





# *Plans for 2004*

---

- ~ *Tribal Harvest*
  - ~ *Creel surveys*
  - ~ *Simple, random, site*
  - ~ *5 days/week*

- ~ *Mark-Recapture Study*
  - ~ *Estimate escapement*
  - ~ *Tag retention*
  - ~ *Average movement rate*

# Conclusions

## *Objective 1*

### *Larval lamprey & Habitat*

- ~ Lamprey were present in 4 of the 13 streams sampled*
- ~ Complex habitat needs including depositional areas and woody debris*
- ~ Small sample sizes made it difficult to find relationships with lamprey presence*





# Conclusions

---

## Objective 2 Timing

### Shitike

~ Peak movement: 2002 – March  
2003 – December

### WSR

~ Peak movement: 2002  
2003

### Ammocoetes

March  
May

### Macrophthalmia

March  
December



# Conclusions

## *Objective 3 Tribal Harvest & Upstream migration*

### Adult Escapement Estimate

- ~ Marked 199 adult lamprey
  - ~ Recaptured 35
  - ~ Tag retention 77%

### Tribal Harvest

- ~ 21 creel interview conducted – 614 lamprey
- ~ Estimated harvest 1,017



# Questions?



*Ammocoete*

*We would like to thank Bonneville Power Administration  
for funding this project.*

*Macrophthalmia*

